



Calhoun: The NPS Institutional Archive
DSpace Repository

Reports and Technical Reports

All Technical Reports Collection

1978-06

Simulation of Nonhomogeneous Poisson Processes by Thinning

Lewis, P.A.W.; Shedler, G.S.

Monterey, California. Naval Postgraduate School

<http://hdl.handle.net/10945/63247>

This publication is a work of the U.S. Government as defined in Title 17, United States Code, Section 101. Copyright protection is not available for this work in the United States.

Downloaded from NPS Archive: Calhoun



<http://www.nps.edu/library>

Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.


Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

LEVEL

②

NPS-55-78-014

NAVAL POSTGRADUATE SCHOOL
Monterey, California



SIMULATION OF
NONHOMOGENEOUS POISSON PROCESSES
BY THINNING

by
P. A. W. Lewis
and
G. S. Shedler

JUNE 1978

Approved for public release; distribution unlimited.

DDC FILE COPY AD A059904

NAVAL POSTGRADUATE SCHOOL
MONTEREY, CALIFORNIA

Rear Admiral Tyler F. Dedman
Superintendent

Jack R. Borsting
Provost

Reproduction of all or part of this report is authorized.

This report was prepared by:

Peter A. W. Lewis
Peter A. W. Lewis, Professor
Department of Operations Research

G.S. Shedler by P.A.W.L.
Gerald S. Shedler
IBM Research Laboratory

Reviewed by:

Released by:

Michael G. Sovereign, Acting
Michael G. Sovereign, Chairman
Department of Operations Research

William M. Tolles
William M. Tolles
Acting Dean of Research

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NPS55-78-14	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Simulation of Nonhomogeneous Poisson Processes by Thinning		5. TYPE OF REPORT & PERIOD COVERED Technical rept
7. AUTHOR(s) Peter A. W. Lewis and G. S. Shedler Gerald S. Shedler		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School Monterey, Ca. 93940		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61153N, RRO14-05-01 N 0001478WR80035
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) RRO1405		12. REPORT DATE June 1978
17. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 43 p.		13. NUMBER OF PAGES 27
18. SUPPLEMENTARY NOTES		15. SECURITY CLASS. (of this report) Unclassified
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non-homogeneous Poisson processes, two-dimensional nonhomogeneous Poisson processes, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics.		16a. DECLASSIFICATION/DOWNGRADING SCHEDULE
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional and two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need for numerical integration of the rate function, for ordering of points, and for generation of Poisson variates.		

SIMULATION OF NONHOMOGENEOUS POISSON PROCESSES BY THINNING

by

P. A. W. Lewis*
Naval Postgraduate School
Monterey, CA 93940

G. S. Shedler
IBM Research Laboratory
San Jose, CA 95193

JUNE 1978

ADDRESS _____
CITY _____ STATE _____ ZIP _____
COUNTRY _____
NAME _____
BY _____
DISTRIBUTIONAL ACTIVITY CODES _____

A

* Support under Office of Naval Research Grant NR-42-343 is gratefully acknowledged.

SIMULATION OF NONHOMOGENEOUS POISSON PROCESSES BY THINNING

by

P. A. W. Lewis
Naval Postgraduate School
Monterey, CA 93940

and

G. S. Shedler
IBM Research Laboratory
San Jose, CA 95193

ABSTRACT

A simple and relatively efficient method for simulating one-dimensional and two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need for numerical integration of the rate function, for ordering of points, and for generation of Poisson variates.

KEYWORDS: Poisson processes, nonhomogeneous Poisson processes, one-dimensional nonhomogeneous Poisson processes, two-dimensional nonhomogeneous Poisson processes, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics.

1. INTRODUCTION

The one-dimensional nonhomogeneous Poisson process (see e.g., Cox and Lewis, 1966, pp. 28-29; Çinlar, 1975, pp. 94-101) has the characteristic properties that the numbers of points in any finite set of nonoverlapping intervals are mutually independent random variables, and that the number of points in any interval has a Poisson distribution. The most general nonhomogeneous Poisson process can be defined in terms of a monotone nondecreasing right-continuous function $\Lambda(x)$ which is bounded in any finite interval. Then the number of points in any finite interval, for example $(0, x_0]$, has a Poisson distribution with parameter $\mu_0 = \Lambda(x_0) - \Lambda(0)$. In this paper it is assumed that $\Lambda(x)$ is continuous, but not necessarily absolutely continuous. The right derivative $\lambda(x)$ of $\Lambda(x)$ is called the rate function of the process; $\Lambda(x)$ is called the integrated rate function and has the interpretation that for $x \geq 0$, $\Lambda(x) - \Lambda(0) = E[N(x)]$, where $N(x)$ is the total number of points in $(0, x]$. Note that $\lambda(x)$ may jump at points at which $\Lambda(x)$ is not absolutely continuous. In contrast to the homogeneous Poisson process, i.e., $\lambda(x)$ a constant (usually denoted by λ), the intervals between the points in a one-dimensional nonhomogeneous Poisson process are neither independent nor identically distributed.

Applications of the one-dimensional nonhomogeneous Poisson process include modelling of the incidence of coal-mining disasters (Cox and Lewis, 1966), the arrivals at an intensive care unit (Lewis, 1972), transaction processing in a data base management system (Lewis and Shedler, 1976b), occurrences of major

freezes in Lake Constance (Steinijans, 1976), and geomagnetic reversal data (Reyment, 1976). The statistical analysis of trends in a one-dimensional nonhomogeneous Poisson process, based on the assumption of an exponential polynomial rate function, is discussed by Cox and Lewis (1966); Cox (1972), Lewis (1972), and Lewis and Shedler (1976b).

There are a number of methods for simulating nonhomogeneous Poisson process which we review briefly.

(i) Time-scale transformation of a homogeneous (rate one)

Poisson process via the inverse of the (continuous) integrated rate function $\Lambda(x)$ constitutes a general method for generation of the nonhomogeneous Poisson process (cf., Çinlar, 1975 pp. 96-97). This method is based on the result that X_1, X_2, \dots are the points in a nonhomogeneous Poisson process with continuous integrated rate function $\Lambda(x)$ if and only if $X'_1 = \Lambda(X_1)$, $X'_2 = \Lambda(X_2)$, ... , are the points in a homogeneous Poisson process of rate one. The time-scale transformation method is a direct analogue of the inverse probability integral transformation method for generating (continuous) nonuniform random numbers. For many rate functions, inversion of $\Lambda(x)$ is not simple and must be done numerically; cf., Gilchrist (1977) and Patrow (1977). The resulting algorithm for generation of the nonhomogeneous Poisson process may be far less efficient than generation based on other methods; see e.g., Lewis and Shedler (1976a, 1977) and Patrow (1977) for discussion of special methods for efficiently generating the nonhomogeneous Poisson process with log-linear and log-quadratic rate functions.

- (ii) A second general method for generating a nonhomogeneous Poisson process with integrated rate function $\Lambda(x)$ is to generate the intervals between points individually, an approach which may seem more natural in the event-scheduling approach to simulation. Thus, given the points $X_1 = x_1, X_2 = x_2, \dots, X_i = x_i$, with $x_1 < x_2 < \dots < x_i$, the interval to the next point, $X_{i+1} - X_i$, is independent of x_1, \dots, x_{i-1} and has distribution function $F(x) = 1 - \exp[-\{\Lambda(x_i + x) - \Lambda(x_i)\}]$. It is possible to find the inverse distribution function $F^{-1}(\cdot)$, usually numerically, and generate $X_{i+1} - X_i$ according to $X_{i+1} - X_i = F^{-1}(U_i)$, where U_i is a uniform random number on the interval $(0,1)$. Note, however, that this not only involves computing the inverse distribution function for each interval $X_{i+1} - X_i$, but that each distribution has different parameters and possibly a different form. An additional complication is that $X_{i+1} - X_i$ is not necessarily a proper random variable, i.e., there may be positive probability that $X_{i+1} - X_i$ is infinite. It is necessary to take this into account for each interval $X_{i+1} - X_i$ before the inverse probability integral transformation is applied. The method is therefore very inefficient with respect to speed, more so than the time-scale transformation method.
- (iii) In a third method, simulation of a nonhomogeneous Poisson process in a fixed interval $(0, x_0]$ can be reduced to the generation of a Poisson number of order statistics from a fixed density function by the following result (cf., Cox and Lewis, 1966, p. 45). If X_1, X_2, \dots, X_n are the points of the

nonhomogeneous Poisson process in $(0, x_0]$, and if $N(x_0) = n$, then conditional on having observed $n(> 0)$ points in $(0, x_0]$, the X_i are distributed as the order statistics from a sample of size n from the distribution function $\{\Lambda(x) - \Lambda(0)\} / \{\Lambda(x_0) - \Lambda(0)\}$, defined for $0 < x \leq x_0$. Generation of the nonhomogeneous Poisson process based on order statistics is in general more efficient (with respect to speed) than either of the previous two methods. Of course, a price is paid for this greater efficiency. First, it is necessary to be able to generate Poisson variates, and second, more memory is needed than in the interval-by-interval method in order to store the sequence of points. Enough memory must be provided so that with very high probability the random number of points generated in the interval can be stored. Recall that the number of points in the interval $(0, x_0]$ has a Poisson distribution with mean $\mu_0 = \Lambda(x_0) - \Lambda(0)$. Memory of size, e.g., $\mu_0 + 4\mu_0^{1/2}$ will ensure that overflow will occur on the average in only one out of approximately every 40,000 realizations. This probability is small enough so that in the case of overflow, the realization of the process can generally be discarded.

- (iv) Again, there is a very particular and very efficient method for simulation of nonhomogeneous Poisson processes with log-linear rate function (Lewis and Shedler, 1976a) which, at the cost of programming complexity and memory, can be used to obtain an efficient simulation method for other rate function, as in Lewis and Shedler (1977).

In this paper a new method is given for simulating a non-homogeneous Poisson process which is not only conceptually simple, but is also computationally simple and relatively efficient. In fact, at the cost of some efficiency, the method can be applied to simulate the given nonhomogeneous Poisson process without the need for numerical integration or routines for generating Poisson variates. Used in conjunction with the special methods given by Lewis and Shedler (1976a, 1977), the method can be used to generate quite efficiently nonhomogeneous Poisson processes with rather complex rate function, in particular combinations of long-term trends and fixed-cycle effects. The method is also easily extended to the problem of generating the two-dimensional nonhomogeneous Poisson process.

2. SIMULATION OF ONE-DIMENSIONAL NONHOMOGENEOUS POISSON PROCESSES

Simulation of a nonhomogeneous Poisson process with general rate function $\lambda(x)$ in a fixed interval can be based on thinning of a nonhomogeneous Poisson process with rate function $\lambda^*(x) \geq \lambda(x)$. The basic result is

THEOREM 1. Consider a one-dimensional nonhomogeneous Poisson process $\{N^*(x): x \geq 0\}$ with rate function $\lambda^*(x)$, so that the number of points, $N^*(x_0)$, in a fixed interval $(0, x_0]$ has a Poisson distribution with parameter $\mu_0^* = \Lambda^*(x_0) - \Lambda^*(0)$. Let $x_1^*, x_2^*, \dots, x_{N^*(x_0)}^*$ be the points of the process in the interval

$(0, x_0]$. Suppose that for $0 \leq x \leq x_0$, $\lambda(x) \leq \lambda^*(x)$. For $i = 1, 2, \dots, n$, delete the point X_i^* with probability $1 - \lambda(X_i^*)/\lambda^*(X_i^*)$; then the remaining points form a nonhomogeneous Poisson process $\{N(x): x \geq 0\}$ with rate function $\lambda(x)$ in the interval $(0, x_0]$.

Proof. Since $\{N^*(x): x \geq 0\}$ is a nonhomogeneous Poisson process, and points are deleted independently, it is clear that the number of points in $\{N(x): x \geq 0\}$ in any set of nonoverlapping intervals are mutually independent random variables. Thus, it is sufficient to show that the number of points $N(a, b)$ in $\{N(x): x \geq 0\}$ in an arbitrary interval $(a, b]$ with $0 \leq a < b \leq x_0$ has a Poisson distribution with parameter $\Lambda(b) - \Lambda(a)$. Observe that with $p(a, b) = \{\Lambda(b) - \Lambda(a)\}/\{\Lambda^*(b) - \Lambda^*(a)\}$, we have the conditional probability

$$P\{N(a, b) = n | N^*(a, b) = k\} = \begin{cases} 1 & \text{if } n = k = 0 \\ \{p(a, b)\}^n \{1 - p(a, b)\}^{k-n} & \text{if } k \geq n \geq 0 \\ & \text{and } k \geq 1 \\ 0 & \text{if } n \geq 1 \\ & \text{and } k < n. \end{cases} \quad (1)$$

Equation (1) is a consequence of the well-known result that, conditional on n (>0) points in the interval $(a, b]$, the joint density of the n points in the process $\{N^*(x): x \geq 0\}$ is

$\lambda^*(x_1) \cdots \lambda^*(x_n) / \{\lambda^*(b) - \lambda^*(a)\}^n$. The desired result is obtained in a straightforward manner from Equation (1) by removing the condition.

Theorem 1 is the basis for the method of simulating non-homogeneous Poisson processes given in this paper.

ALGORITHM 1. One-dimensional nonhomogeneous Poisson process.

1. Generate points in the nonhomogeneous Poisson process $\{N^*(x)\}$ with rate function $\lambda^*(x)$ in the fixed interval $(0, x_0]$. If the number of points generated, n^* , is such that $n^* = 0$, exit; there are no points in the process $\{N(x)\}$.
2. Denote the (ordered) points by $X_1^*, X_2^*, \dots, X_{n^*}^*$. Set $i = 1$ and $k = 0$.
3. Generate U_i , uniformly distributed between 0 and 1. If $U_i \leq \lambda(X_i^*) / \lambda^*(X_i^*)$, set k equal to $k+1$ and $X_k = X_i^*$.
4. Set i equal to $i+1$. If $i \leq n^*$, go to 3.
5. Return X_1, X_2, \dots, X_n , where $n = k$, and also n .

In the case where

- (i) $\{N^*(x)\}$ is a homogeneous Poisson process with $\lambda^*(x) = \lambda^*$;
- (ii) the minimum of $\lambda(x)$, say $\underline{\lambda}$, is known, and
- (iii) generation of uniformly distributed variates is computationally costly,

considerable speedup can be obtained by noting that X_i^* is always accepted if $U_i \leq \underline{\lambda} / \lambda^*$. This obviates, in some cases, computation

of $\lambda(x)$, which is the main source of inefficiency in the algorithm. Moreover, in this case $\lambda^* U_i / \lambda$ can be used as the next uniformly distributed variate.

The method of thinning in this simple form, i.e., $\lambda^*(x) = \lambda^* \geq \max_{0 \leq x \leq x_0} \lambda(x)$, can also be used to provide an algorithm for generating a nonhomogeneous Poisson process on an interval-by-interval basis, as discussed in subsection (ii) of Section 1. The interval to the next point $X_{i+1} - X_i$ is obtained by generating and cumulating exponential(λ^*) random numbers E_1^*, E_2^*, \dots , until for the first time $U_j \leq \lambda(X_1 + E_1^* + \dots + E_j^*) / \lambda^*$, where the U_j are independent uniform random numbers between 0 and 1. This algorithm is considerably simpler than the interval-by-interval algorithm of Section 1 since it requires no numerical integration, only the availability of uniform random numbers.

3. DISCUSSION OF THE METHOD OF THINNING

(i) Relationship to acceptance-rejection method

The method of thinning of Algorithm 1 is essentially the obverse of the conditional method of Section 1, using conditioning and acceptance-rejection techniques to generate the random variables with density function $\lambda(x)/\{\Lambda(x) - \Lambda(0)\}$ (Lewis and Shedler, 1977, Algorithm 3). The differences are subtle, but computationally important. In the acceptance-rejection method, it is first necessary to generate a Poisson variate with mean $\mu_0 = \Lambda(x_0) - \Lambda(0)$, and this involves an integration of the rate function $\lambda(x)$. Then the Poisson (μ_0) number, n , of variates generated by acceptance-rejection must be ordered to give X_1, X_2, \dots, X_n .

(ii) Simplest form of the thinning algorithm

In the simplest form of the method of thinning, $\lambda^*(x)$ is taken to be a constant λ^* , so that, for instance, the points $X_1^*, X_2^*, \dots, X_{n^*}^*$ can be generated by cumulating exponential (λ^*) variates until the sum is greater than x_0 (cf., Lewis and Shedler, 1976a, Algorithm 1). Thinning is then applied to the generated points. No ordering, no integration of $\lambda(x)$ and no generator of Poisson variates is required. Of course for both algorithms to be efficient, computation of $\lambda(x)$ and $\lambda^*(x)$ must be easy relative to computation of the inverse of $\Lambda(x)$.

(iii) Efficiency

For the thinning algorithm (as well as the algorithm based on conditioning and acceptance-rejection) efficiency, as measured by the number of points deleted, is proportional to $\mu_0/\mu_0^* = \{\Lambda(x_0) - \Lambda(0)\}/\{\Lambda^*(x_0) - \Lambda^*(0)\}$; this is the ratio of the areas between 0 and x_0 under $\lambda(x)$ and $\lambda^*(x)$. Thus, $\lambda^*(x)$ should be as close as possible to $\lambda(x)$ consistent with ease of generating the nonhomogeneous Poisson process $\{N^*(x): x \geq 0\}$.

(iv) An example: fixed cycle plus trend

To illustrate the applicability of the thinning algorithm, consider its use in conjunction with the algorithms given by Lewis and Shedler (1976a, 1977) for log-linear and log-quadratic rate functions. Assume that it is necessary to simulate a nonhomogeneous Poisson process whose rate function increases quadratically with time but also has a fixed-period cycle, e.g.,

$$\lambda(x) = \exp\{\alpha_0 + \alpha_1 x + \alpha_2 x^2 + K \sin(\omega_0 x + \theta)\},$$

$$0 \leq x \leq x_0; \quad K \geq 0; \quad 0 < \theta \leq 2\pi; \quad \omega_0 > 0.$$

This is the model found by Lewis (1972) for arrivals at an intensive care unit, where there is a strong time-of-day effect. Thus if $\omega_0 = 2\pi/T_0$, then the period $T_0 = 1$ day. Computation of $\Lambda^{-1}(\cdot)$ is difficult. To determine $\lambda^*(x)$, note that

$$\lambda(x) \leq \lambda^*(x) = \exp\{\alpha_0 + K + \alpha_1 x + \alpha_2 x^2\},$$

and therefore

$$\lambda(x)/\lambda^*(x) = \exp[K\{1 - \sin(\omega_0 x + \theta)\}] .$$

Thus in Step 3 of Algorithm 1, U_i is compared to $\exp[K\{1 - \sin(\omega_0 X_i^* + \theta)\}]$. Equivalently, if unit exponential variates E_i are available, it is faster to compare E_i to $K\{1 - \sin(\omega_0 X_i^* + \theta)\}$, accepting X_i^* if $E_i > K\{1 - \sin(\omega_0 X_i^* + \theta)\}$.

The main computational expense here is generation of the E_i and computation of the sine function, both n^* times. The expense involved in computation of the sine function can be reduced by noting that the point X_i^* is always accepted if E_i is greater than $2K$. This will be a great saving if the cyclic effect is minor (K small). The number of E_i generated can be reduced by noting that if, in one step of the algorithm, E_i is observed to be greater than δ , then $E_i^* = E_i - \delta$ can be used as an (independent) unit exponential variate in the next step. The above procedure can be extended to the case of a trend with two fixed-period cycles, e.g., a time-of-day and a time-of-week effect.

4. SIMULATION OF TWO-DIMENSIONAL HOMOGENEOUS POISSON PROCESSES

The two-dimensional homogeneous Poisson process (of rate $\lambda > 0$) is defined by the properties that the numbers of points in any finite set of nonoverlapping regions having areas in the usual geometric sense are mutually independent, and that the number of points in any region of area A has a Poisson distribution with mean λA ;

see, e.g., Karlin and Taylor (1975), pp. 31-32. Note that the number of points in a region R depends on its area, but not on its shape or location. The homogeneous Poisson process arises as a limiting two-dimensional point process with respect to a number of limiting operations; cf., Goldman (1967a,b). Properties of the process are given by Miles (1970). Applications of the two-dimensional homogeneous Poisson process to problems in ecology and forestry have been discussed by Thompson (1955) and Holgate (1972). The model also arises in connection with naval search and detection problems.

In considering the two-dimensional homogeneous Poisson process, projection properties of the process depend quite critically on the geometry of the regions considered. These projection properties are simple for rectangular and circular regions, and make simulation of the homogeneous process quite easy. We consider these two cases separately.

(i) Homogeneous Poisson Processes in a Rectangle

The following two theorems form the basis for simulation of the two-dimensional homogeneous Poisson process in a rectangle.

THEOREM 2. Consider a two-dimensional homogeneous Poisson process of rate λ , so that the number of points in a fixed rectangle $R = \{(x,y): 0 < x \leq x_0, 0 < y \leq y_0\}$ has a Poisson distribution with parameter $\lambda x_0 y_0$. If $(X_1, Y_1), (X_2, Y_2), \dots, (X_N, Y_N)$ denote the position of the points of the process in R , labelled so that

$x_1 < x_2 < \dots < x_N$, then x_1, x_2, \dots, x_N form a one-dimensional homogeneous Poisson process on $0 < x \leq x_0$ of rate λy_0 . If the points are relabelled $(x'_1, y'_1), (x'_2, y'_2), \dots, (x'_N, y'_N)$ so that $y'_1 < y'_2 < \dots < y'_N$, then y'_1, y'_2, \dots, y'_N form a one-dimensional homogeneous Poisson process on $0 < y \leq y_0$ of rate λx_0 .

Proof. The number of points in an interval on the x-axis, say, $(a, b]$ is the number of points in the rectangle bounded by the lines $x = a, x = b, y = 0$, and $y = y_0$. This number is therefore independent of the number of points in any similar nonoverlapping rectangle bounded on the x-axis by $x = a', x = b'$, i.e., the number of points in the interval $(a', b']$. This establishes the independent increment property for a one-dimensional Poisson process. The Poisson distribution of the number of points in $(a, b]$ follows from the fact that it is equal to the number of points in the rectangle bounded by $x = a, x = b, y = 0$, and $y = y_0$, and the latter has a Poisson distribution with parameter $\lambda y_0 (b-a)$. An analogous argument shows that the process formed on the y-axis by y'_1, y'_2, \dots, y'_N is Poisson.

Conditional properties of the Poisson process in a rectangle are established next. The important thing to note is that while the processes obtained by projection of the points onto the x and y axes are not independent, there is a type of conditional independence which makes it easy to simulate the two-dimensional process.

THEOREM 3. Assume that a two-dimensional homogeneous Poisson process of rate λ is observed in a fixed rectangle $R = \{(x,y): 0 < x \leq x_0, 0 < y \leq y_0\}$, so that the number of points in R , $N(R)$, has a Poisson distribution with parameter $\lambda x_0 y_0$. If $N(R) = n > 0$ and if $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$ denote the points, labelled so that $X_1 < X_2 < \dots < X_n$, then conditional on having observed n points in R , the X_1, X_2, \dots, X_n are uniform order statistics on $0 < x \leq x_0$, and Y_1, Y_2, \dots, Y_n are independent and uniformly distributed on $0 < y \leq y_0$, independent of the X_i .

Proof. If there are N points in the rectangle, form N vertical strips, from 0 to y_0 and from X_i to $X_i + dx_i$, such that each strip contains only one of the N points. The position of Y_i on the vertical line through X_i is that of an event in a Poisson process of rate λdx_i , given that only one event occurs. But this means that Y_i is uniformly distributed between 0 and y_0 . Moreover, this is true irrespective of where X_i occurs; therefore Y_i is independent of X_i . Also, occurrences in all N strips are independent, and therefore Y_i is independent of the other Y_j and X_j positions, $j \neq i$. Thus the Y_i are a random sample of size N from a uniform $(0, y_0)$ distribution, independent of the X_i . Now condition on $N = n (> 0)$; since by Theorem 2 the X_i form a Poisson process they are, by well-known results, order statistics from a uniform $(0, x_0)$ sample and are independent of the fixed size Y_i population; thus the pairs (X_i, Y_i) are mutually independent.

COROLLARY. Denote the Poisson points by $(X_1, Y_1), (X_2, Y_2), \dots$, where the index does not necessarily denote an ordering on either axis. Conditionally, the pairs $(X_1, Y_1), \dots, (X_N, Y_N)$ are independent random variables. Furthermore, for each pair, X_i is distributed uniformly between 0 and x_0 , independently of Y_i , which is uniformly distributed between 0 and y_0 .

From the two theorems, the following simulation procedure is obtained.

ALGORITHM 2. Two-dimensional homogeneous Poisson process in a rectangle.

1. Generate points in the one-dimensional homogeneous Poisson process of rate λy_0 on $(0, x_0]$. If the number of points generated, n , is such that $n = 0$, exit; there are no points in the rectangle.
2. Denote the points generated by $\lambda_1 < X_2 < \dots < X_n$.
3. Generate Y_1, Y_2, \dots, Y_n as independent, uniformly distributed random numbers on $(0, y_0]$.
4. Return $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$ as the coordinates of the two-dimensional homogeneous Poisson process in the rectangle, and n .

Note that generation of the points X_1, X_2, \dots, X_N in Steps 1 and 2 can be accomplished by cumulating exponential(λy_0) random numbers. Alternatively, after generating a Poisson random number $N = n$ (with parameter $\lambda x_0 y_0$), n independent, uniformly distributed random numbers on $(0, x_0]$ can be ordered; see Lewis and Shedler (1976a), p. 502.

Another algorithm for generation of the two-dimensional Poisson process in a rectangle can be based on the Corollary to Theorem 3.

(ii) Homogeneous Poisson Processes in a Circle

The following two theorems form the basis for simulation of the two-dimensional homogeneous Poisson process in a fixed circle of radius r_0 .

Fix the origin and initial line of polar coordinates r and θ so that the origin is the center of the circle and the initial line is horizontal. We consider the projection of the points (R_i, θ_i) , of the Poisson process circularly onto the r -axis (R_i) and radially onto the circumferential θ -axis (θ_i) . The number of points projected onto the r -axis in the interval $(0, r]$, where $r \leq r_0$, is the number of points in the circle of radius r and area πr^2 ; thus the number of points in $(0, r]$ has a Poisson distribution with parameter $\lambda \pi r^2$. Consequently, if the projection process on the r -axis is a Poisson process, it must have integrated rate function $\Lambda(r) = \lambda \pi r^2$, with $\Lambda(0) = 0$.

Similarly, the number of points on the circumferential arc of the fixed circle (radius r_0) from 0 to θ is the number of points in the sector of the circle defined by radial lines at angles 0 and θ ; thus, the number of points on the arc from 0 to θ has a Poisson distribution with parameter $\lambda \pi r_0^2 \times \frac{\theta}{2\pi} = \theta \lambda r_0^2 / 2$. Accordingly, if the projection process on the θ -axis is a Poisson process, it must have integrated rate function $\Lambda(\theta) = \theta \lambda r_0^2 / 2$, with $\Lambda(0) = 0$.

We now assert that the projection processes are in fact Poisson processes. Since proofs of these theorems are directly analogous to the proofs of Theorems 2 and 3, they are omitted.

THEOREM 4. Consider a two-dimensional homogeneous Poisson process of rate λ so that the number N of points in a fixed circular area C of radius r_0 and area πr_0^2 has a Poisson distribution with parameter $\lambda \pi r_0^2$. If $(R_1, \theta_1), (R_2, \theta_2), \dots, (R_N, \theta_N)$ denote the points of the process in C , labelled so that $R_1 < R_2 < \dots < R_N$, then R_1, R_2, \dots, R_N form a one-dimensional nonhomogeneous Poisson process on $0 \leq r \leq r_0$ with rate function $\lambda(r) = 2\pi\lambda r$. If the points are relabelled $(R'_1, \theta'_1), (R'_2, \theta'_2), \dots, (R'_N, \theta'_N)$ so that $\theta'_1 < \theta'_2 < \dots < \theta'_N$, then $\theta'_1, \theta'_2, \dots, \theta'_N$ form a one-dimensional homogeneous Poisson process on $0 < \theta \leq 2\pi$ of rate $\lambda r_0^2/2$.

THEOREM 5. Assume that a two-dimensional Poisson process of rate λ is observed in a fixed circular area C of radius r_0 so that the number of points in C , $N(C)$, has a Poisson distribution with parameter $\lambda \pi r_0^2$. If $N(C) = n > 0$ and if $(R_1, \theta_1), (R_2, \theta_2), \dots, (R_n, \theta_n)$ with $R_1 < R_2 < \dots < R_n$ denote the points, then conditional on having observed n points in C , the R_1, R_2, \dots, R_n are order statistics from the density $f(r) = 2r/r_0^2$ concentrated on $0 \leq r \leq r_0$, and $\theta_1, \theta_2, \dots, \theta_n$ are independent and uniformly distributed on $0 < \theta \leq 2\pi$, independent of the R_i .

These theorems lead to the following simulation procedure.

ALGORITHM 3. Two-dimensional homogeneous Poisson process in a circular area.

1. Generate n as a Poisson random number with parameter $\lambda\pi r_0^2$. If $n = 0$, exit; there are no points in C .
2. Generate n independent random numbers having density function $f(r) = 2r/r_0^2$ and order to obtain $R_1 < R_2 < \dots < R_n$.
3. Generate $\theta_1, \theta_2, \dots, \theta_n$ independent, uniformly distributed random numbers on $(0, 2\pi]$.
4. Return $(R_1, \theta_1), (R_2, \theta_2), \dots, (R_n, \theta_n)$, and n .

Note that the wedge-shaped density $2r/r_0^2$ can be generated by scaling the maximum of two independent uniform $(0,1)$ random numbers.

Direct generation of homogeneous Poisson points in non-circular or non-rectangular regions is difficult. The processes obtained by projection of the points on the two axes are nonhomogeneous Poisson processes with complex rate functions determined by the geometry of the region. However, the conditional independence which is found in circular and rectangular regions (Theorems 3 and 5) for the processes on the two axes is not present. In particular, given that there are n points $(X_1, Y_1), \dots, (X_n, Y_n)$ in a non-rectangular region, the pairs (X_i, Y_i) are mutually independent, but X_i is in general not independent of Y_i , $i = 1, \dots, n$. Therefore, it is simpler to enclose the

region in either a circle or a rectangle, generate a homogeneous Poisson process in the enlarged area, and subsequently exclude points outside of the given region.

5. SIMULATION OF TWO-DIMENSIONAL NONHOMOGENEOUS POISSON PROCESSES

The two-dimensional nonhomogeneous Poisson process $\{N(x,y): x \geq 0, y \geq 0\}$ is specified by a positive rate function $\lambda(x,y)$ which for simplicity is assumed here to be continuous. Then the process has the characteristic properties that the numbers of points in any finite set of nonoverlapping regions having areas in the usual geometric sense are mutually independent, and that the number of points in any such region R has a Poisson distribution with mean $\Lambda(R)$; here $\Lambda(R)$ denotes the integral of $\lambda(x,y)$ over R , i.e., over the entire area of R .

Applications of the two-dimensional nonhomogeneous Poisson process include problems in forestry and naval search and detection. The use of the process as a model for the pattern of access to the storage subsystem of a computer system will be reported elsewhere. Detection and statistical analysis of trends in the two-dimensional nonhomogeneous Poisson process is discussed by Rantschler (1973).

Theorem 1 dealing with thinning of one-dimensional nonhomogeneous Poisson processes generalizes to two-dimensional nonhomogeneous Poisson processes. Thus, suppose that

$\lambda(x,y) \leq \lambda^*(x,y)$ in a fixed rectangular region of the plane.

If a nonhomogeneous Poisson process with rate function $\lambda^*(x,y)$ is thinned according to $\lambda(x,y)/\lambda^*(x,y)$ (i.e., each point

(X_i, Y_i) is deleted independently if a uniform $(0,1)$ random number U_i is greater than $\lambda(X_i, Y_i)/\lambda^*(X_i, Y_i)$, the result is a nonhomogeneous Poisson process with rate function $\lambda(x, y)$. The proof is a direct analogue of the proof for the one-dimensional case.

The nonhomogeneous Poisson process with rate function $\lambda(x, y)$ in an arbitrary but fixed region R can be generated by enclosing the region R either in a rectangle or a circle, and applying Algorithm 2 or Algorithm 3. The following procedure assumes that the region R has been enclosed in a rectangle R^* , and that $\lambda^* = \max\{\lambda(x, y) : x, y \in R\}$ has been determined; here the bounding process is homogeneous with rate λ^* in the rectangle R^* .

ALGORITHM 4. Two-dimensional nonhomogeneous Poisson process.

1. Using Algorithm 2, generate points in the homogeneous Poisson process of rate λ^* in the rectangle R^* . If the number of points, n^* , is such that $n^* = 0$, exit; there are no points in the nonhomogeneous Poisson process.
2. From the n^* points generated in 1, delete the points that are not in R , and denote the remaining points by $(X_1^*, Y_1^*), (X_2^*, Y_2^*), \dots, (X_m^*, Y_m^*)$ with $X_1^* < X_2^* < \dots < X_m^*$. Set $i = 1$ and $k = 0$.
3. Generate U_i uniformly distributed between 0 and 1. If $U_i \leq \lambda(X_i^*, Y_i^*)/\lambda^*$, set $k = k+1$, $X_k = X_i^*$ and $Y_k = Y_i^*$.
4. Set i equal to $i+1$. If $i \leq m^*$, go to 3.
5. Return $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$, where $n = k$, and n .

It is not necessary that the bounding process have a constant rate λ^* . Theorems 2 and 4 can be generalized to certain cases where the process is nonhomogeneous (cf., Bartlett, 1974), for instance $\lambda(x,y) = \rho(x) \psi(y)$. Thus, a tighter bounding process which is nonhomogeneous may possibly be obtained. It is not simple to see how much efficiency could be gained by doing this, as opposed to using a two-dimensional homogeneous Poisson process for the bounding process. Again, as in the one-dimensional case, savings in computing $\lambda(x,y)$ can be obtained by computing its minimum beforehand, and the U_i 's can be reused by scaling.

6. COMPARISONS AND CONCLUDING REMARKS

The method of thinning presented in this paper for simulating one-dimensional and two-dimensional nonhomogeneous Poisson processes with given rate function can be carried out in a computationally simple way by using a bounding process which is homogeneous with a rate function equal to the maximum value of the given rate function. No numerical integration, ordering or generation of Poisson variates is required, only the ability to evaluate the given rate function. The thinning algorithm appears to be particularly attractive in the two-dimensional case where there seem to be no competing algorithms.

The thinning algorithm can also be implemented more efficiently at the cost of programming complexity and by using a nonhomogeneous bounding process. In particular the method can be used in conjunction with the special algorithms given by Lewis and Shedler (1976a, 1977).

It is also possible to extend the method of thinning to simulation of doubly stochastic or conditioned Poisson processes. This will be discussed elsewhere.

REFERENCES

- Ahrens, J. and Dieter, U. (1974a). Computer methods for sampling from gamma, beta, Poisson and binomial distributions. Computing 12, 223-46.
- Ahrens, J. and Dieter, U. (1974b). Non Uniform Random Numbers. Graz, Austria: Technische Hochschule in Graz.
- Bartlett, M.S. (1964). The spectral analysis of two-dimensional point processes. Biometrika 51, 299-311.
- Cinlar, E. (1975). Introduction to Stochastic Processes. Englewood Cliffs, New Jersey: Prentice-Hall.
- Cox, D.R. The statistical analysis of dependencies in point processes. In Stochastic Point Processes, Ed. P.A.W. Lewis, pp. 55-56. New York: Wiley.
- Cox, D.R. and Lewis, P.A.W. (1966). The Statistical Analysis of Series of Events. London: Methuen.
- Gilchrist, R. (1977). Some problems encountered in generating realisations from a useful non-homogeneous Poisson process. In Proceedings of the European Congress of Statisticians 1976, Ed. J. R. Barra et al. Amsterdam: North Holland.
- Goldman, J.R. (1967a). Stochastic point processes: limit theorems. Ann. Math. Stat. 38, 771-79.
- Goldman, J.R. (1967b). Infinitely divisible point processes in R^n . J. Math. Anal. Appl. 17, 133-46.
- Holgate, P. (1972). The use of distance methods for the analysis of spatial distributions of points. In Stochastic Point Processes, Ed. P.A.W. Lewis, pp. 122-35. New York: Wiley.
- Karlin, S. and Taylor, H.M. (1975). A First Course in Stochastic Processes. New York: Academic Press.
- Lewis, P.A.W. (1972). Recent advances in the statistical analysis of univariate point processes. In Stochastic Point Processes, Ed. P.A.W. Lewis, pp. 1-54. New York: Wiley.
- Lewis, P.A.W. and Shedler, G.S. (1976a). Simulation of non-homogeneous processes with log-linear rate function. Biometrika 63, 501-05.

- Lewis, P.A.W. and Shedler, G.S. (1976b). Statistical analysis of a non-stationary series of events in a data base system. IBM J. Res. Devel. 20, 465-82.
- Lewis, P.A.W. and Shedler, G.S. (1977). Simulation of nonhomogeneous Poisson processes with degree-two exponential polynomial rate function. IBM Research Report RJ 1953. San Jose, California.
- Miles, R.E. (1970). On the homogeneous planar Poisson point process. Mathematical Biosciences 6, 85-127.
- Patrow, M.L. (1977). A comparison of two algorithms for the simulation of non-homogeneous Poisson processes with degree-two exponential polynomial intensity function. M.S. Thesis., Dept. of Operations Research, Naval Postgraduate School. Monterey, California (National Technical Information Service, Springfield, Va.; AD-A047164).
- Rantschler, R.D. (1973). The detection and estimation of trends in two-dimensional Poisson processes. M.S. Thesis, Dept. of Operations Research, Naval Postgraduate School, Monterey, California (National Technical Information Service, Springfield, Va.; AD-A76136).
- Reyment, R.A. (1976). Geomagnetic reversal data statistically appraised. Tectonophysics 31, 73-91.
- Steinijans, V. (1976). A stochastic point process model of the occurrence of major freezes in Lake Constance. Appl. Statistics 25, 58-61.
- Thompson, H.R. (1955). Spatial point processes with applications to ecology. Biometrika 42, 102-15.
- Vere-Jones, D. (1970). Stochastic models for earthquake occurrence. J. Roy. Statist. Soc. B 32, 1-62.

DISTRIBUTION LIST

NO. OF COPIES

STATISTICS AND PROBABILITY PROGRAM
OFFICE OF NAVAL RESEARCH
CODE 436
ARLINGTON
VA

22217

1

OFFICE OF NAVAL RESEARCH
NEW YORK AREA OFFICE
715 BROADWAY - 5TH FLOOR
ATTN: DR. ROGER GRAFTON
NEW YORK, NY

10003

1

DIRECTOR
OFFICE OF NAVAL RESEARCH BRANCH OFF
536 SOUTH CLARK STREET
ATTN: DEPUTY AND CHIEF SCIENTIST
CHICAGO, IL

60605

1

LIBRARY
NAVAL OCEAN SYSTEMS CENTER
SAN DIEGO
CA

92152

1

NAVY LIBRARY
NATIONAL SPACE TECHNOLOGY LAB
ATTN: NAVY LIBRARIAN
BAY ST. LOUIS
MS

39522

1

NAVAL ELECTRONIC SYSTEMS COMMAND
NAVELEX 320
NATIONAL CENTER NO. 1
ARLINGTON
VA

20360

1

DIRECTOR NAVAL RESEARCH LABORATORY
ATTN: LIBRARY (ONRL)
CODE 2029
WASHINGTON, D.C.

20375

1

DEFENSE DOCUMENTATION CENTER
CAMERON STATION

1

ALEXANDRIA
VIRGINIA 22314

TECHNICAL INFORMATION DIVISION
NAVAL RESEARCH LABORATORY

1

WASHINGTON, D. C.

20375

	NO. OF COPIES
OFFICE OF NAVAL RESEARCH SAN FRANCISCO AREA OFFICE 760 MARKET STREET SAN FRANCISCO CALIFORNIA 94102	1
TECHNICAL LIBRARY NAVAL ORDNANCE STATION INDIAN HEAD MARYLAND 20640	1
NAVAL SHIP ENGINEERING CENTER PHILADELPHIA DIVISION TECHNICAL LIBRARY PHILADELPHIA PENNSYLVANIA 19112	1
BUREAU OF NAVAL PERSONNEL DEPARTMENT OF THE NAVY TECHNICAL LIBRARY WASHINGTON D. C. 20370	1
LIBRARY CODE 3212 NAVAL POSTGRADUATE SCHOOL MONTREY CALIFORNIA 93940	1
PROF. M. AEDEL-HAMEED DEPARTMENT OF MATHEMATICS UNIVERSITY OF NORTH CAROLINA CHARLOTTE NC 28223	1
PROF. T. W. ANDERSON DEPARTMENT OF STATISTICS STANFORD UNIVERSITY STANFORD , CALIFORNIA 94305	1
PROF. F. J. ANSCOMBE DEPARTMENT OF STATISTICS YALE UNIVERSITY NEW HAVEN CONNECTICUT 06520	1
PROF. L. A. ARCIA INSTITUTE OF INDUSTRIAL ADMINISTRATION UNION COLLEGE SCHENECTADY NEW YORK 12308	1
PROF. C. R. BAKER DEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CAROLINA CHAPEL HILL NORTH CAROLINA 27514	1

NO. OF COPIES

PROF. R. E. BECHOFER
DEPARTMENT OF OPERATIONS RESEARCH
CORNELL UNIVERSITY
ITHACA
NEW YORK 14850

1

PROF. N. J. BERSHAD
SCHOOL OF ENGINEERING
UNIVERSITY OF CALIFORNIA
IRVINE,
CALIFORNIA
92664

1

P. J. BICKEL
DEPARTMENT OF STATISTICS
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA

1

94720

PROF. F. W. BLECK
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF PITTSBURGH
PITTSBURGH
PA

1

15260

PROF. JOSEPH BLUM
DEPT. OF MATHEMATICS, STATISTICS
AND COMPUTER SCIENCE
THE AMERICAN UNIVERSITY
WASHINGTON
DC

1

20016

PROF. R. A. BRADLEY
DEPARTMENT OF STATISTICS
FLORIDA STATE UNIVERSITY

1

TALLAHASSEE, FLORIDA 32306

PROF. R. E. BARLOW
OPERATIONS RESEARCH CENTER
COLLEGE OF ENGINEERING
UNIVERSITY OF CALIFORNIA
BERKELEY
CALIFORNIA 94720

1

MR. C. N. BENNETT
NAVAL COASTAL SYSTEMS LABORATORY
CODE P761
PANAMA CITY,
FLORIDA
32401

1

PROF. U. N. BHAT
COMPUTER SCIENCE / OPERATIONS
RESEARCH CENTER
SOUTHERN METHODIST UNIVERSITY
DALLAS
TEXAS 75275

1

PROF. W. R. BLISCHKE
DEPT. OF QUANTITATIVE
BUSINESS ANALYSIS
UNIVERSITY OF SOUTHERN CALIFORNIA
LOS ANGELES, CALIFORNIA

1

90007

NO. OF COPIES

DR. DERRILL J. BORDELON
NAVAL UNDERWATER SYSTEMS CENTER
COCE 21
NEWPORT
RI

1

C2840

J. E. BOYER JR
DEPT. OF STATISTICS
SOUTHERN METHODIST UNIVERSITY
DALLAS
TX

1

75275

DR. J. CHANDRA
U. S. ARMY RESEARCH
P. O. BOX 12211
RESEARCH TRIANGLE PARK,
NORTH CAROLINA
27706

1

PROF. F. CHERNOFF
DEPT. OF MATHEMATICS
MASS INSTITUTE OF TECHNOLOGY
CAMBRIDGE,
MASSACHUSETTS 02139

1

PROF. C. DERMAN
DEPARTMENT OF CIVIL ENGINEERING
AND ENGINEERING MECHANICS
COLUMBIA UNIVERSITY
NEW YORK
NEW YORK

1

10027

PROF. R. L. DISNEY
VIRGINIA POLYTECHNIC INSTITUTE
AND STATE UNIVERSITY
DEPT. OF INDUSTRIAL ENGINEERING
AND OPERATIONS RESEARCH
BLACKSBURG, VA

1

24061

MR. J. DOWLING
DEFENSE LOGISTICS STUDIES
INFORMATION EXCHANGE
ARMY LOGISTICS MANAGEMENT CENTER
FORT LEE
VIRGINIA 20390

1

PROF. J. D. ESARY
DEPT. OF OPERATIONS RESEARCH AND
ADMINISTRATIVE SCIENCES
NAVAL POSTGRADUATE SCHOOL
MONTEREY,
CALIFORNIA 93940

1

DR. M. J. FISCHER
DEFENSE COMMUNICATIONS AGENCY
1860 WIEHLE AVENUE
RESTON
VIRGINIA
22070

1

NO. OF COPIES

PROF. C. P. GAVER
DEPT. OF OPERATIONS RESEARCH
NAVAL POSTGRADUATE SCHOOL
MONTEREY
CA

1

93940

MR. GENE F. GLEISSNER
APPLIED MATHEMATICS LABORATORY
DAVID TAYLOR NAVAL SHIP RESEARCH
AND DEVELOPMENT CENTER
BETHESDA
MD

1

20084

PROF. S. S. GUPTA
DEPARTMENT OF STATISTICS
FURDUE UNIVERSITY
LAFAYETTE
INDIANA 47907

1

PROF. D. L. HANSON
DEPT OF MATH. SCIENCES
STATE UNIVERSITY OF NEW YORK,
BINGHAMTON
BINGHAMTON
NY

1

13901

PROF. F. J. HARRIS
DEPT. OF ELECTRICAL ENGINEERING
SAN DIEGO STATE UNIVERSITY
SAN DIEGO
CA

1

92182

PROF. L. H. HERBACH
DEPT. OF OPERATIONS RESEARCH AND
SYSTEMS ANALYSIS
POLYTECHNIC INSTITUTE OF NEW YORK
BROOKLYN
NY

1

11201

PROF. M. J. HINICH
DEPARTMENT OF ECONOMICS
VIRGINIA POLYTECHNIC INSTITUTE
AND STATE UNIVERSITY
BLACKSBURG,
VIRGINIA 24061

1

PROF. W. M. HIRSCH
INSTITUTE OF MATHEMATICAL SCIENCES
NEW YORK UNIVERSITY
NEW YORK
NEW YORK 10453

1

PROF. C. L. IGLEHART
DEPARTMENT OF OPERATIONS RESEARCH
STANFORD UNIVERSITY
STANFORD,
CALIFORNIA
94350

1

NO. OF COPIES

PROF. J. B. KACANE
DEPARTMENT OF STATISTICS
CARNEGIE-MELLON
PITTSBURGH,
PENNSYLVANIA
15213

1

DR. RICHARD LAU
DIRECTOR
OFFICE OF NAVAL RESEARCH BRANCH OFF
1030 EAST GREEN STREET
PASADENA
CA

1

91101

DR. A. R. LAUFER
DIRECTOR
OFFICE OF NAVAL RESEARCH BRANCH OFF
1030 EAST GREEN STREET
PASADENA
CA

1

91101

PROF. M. LEADBETTER
DEPARTMENT OF STATISTICS
UNIVERSITY OF NORTH CAROLINA
CHAPEL HILL
NORTH CAROLINA 27514

1

DR. J. S. LEE
J. S. LEE ASSOCIATES, INC.
2001 JEFFERSON DAVIS HIGHWAY
SUITE EC2
ARLINGTON
VA

1

22202

PROF. L. C. LEE
DEPARTMENT OF STATISTICS
VIRGINIA POLYTECHNIC INSTITUTE
AND STATE UNIVERSITY
BLACKSBURG
VA

1

24061

PROF. R. S. LEVENWORTH
DEPT. OF INDUSTRIAL AND SYSTEMS
ENGINEERING
UNIVERSITY OF FLORIDA
GAINSVILLE,
FLORIDA 32611

1

PROF. G. LIEBERMAN
STANFORD UNIVERSITY
DEPARTMENT OF OPERATIONS RESEARCH
STANFORD CALIFORNIA 94305

1

NO. OF COPIES

DR. JAMES F. MAAR
NATIONAL SECURITY AGENCY
FORT MEADE, MARYLAND
20755

1

PROF. R. W. MADSEN
DEPARTMENT OF STATISTICS
UNIVERSITY OF MISSOURI
COLUMBIA
MO

1

65201

DR. N. R. MANN
SCIENCE CENTER
ROCKWELL INTERNATIONAL CORPORATION
P.O. BOX 1065
THOUSAND OAKS,
CALIFORNIA 91360

1

DR. W. H. MARLOW
PROGRAM IN LOGISTICS
THE GEORGE WASHINGTON UNIVERSITY
707 22ND STREET, N. W.
WASHINGTON, D. C.
20037

1

PROF. E. MASRY
DEPT. APPLIED PHYSICS AND
INFORMATION SERVICE
UNIVERSITY OF CALIFORNIA
LA JOLLA
CALIFORNIA

1

92093

DR. BRUCE J. McDONALD
SCIENTIFIC DIRECTOR
SCIENTIFIC LIAISON GROUP
OFFICE OF NAVAL RESEARCH
AMERICAN EMBASSY - TOKYO
AFC SAN FRANCISCO

1

96503

PROF. J. A. MUCKSTADT
DEPT. OF OPERATIONS RESEARCH
CORNELL UNIVERSITY
ITHACA,
NEW YORK
14850

1

DR. JANET M. MYRE
THE INSTITUTE OF DECISION SCIENCE
FOR BUSINESS AND PUBLIC POLICY
CLAREMONT MEN'S COLLEGE
CLAREMONT
CA

1

91711

MR. H. NISSELSON
BUREAU OF THE CENSUS
ROOM 2025
FEDERAL BUILDING 3
WASHINGTON,
D. C. 20333

1

NO. OF COPIES

MISS B. S. ORLEANS
NAVAL SEA SYSTEMS COMMAND
(SEA 03F)
RM 10508
ARLINGTON VIRGINIA 20360

1

PROF. C. B. OWEN
DEPARTMENT OF STATISTICS
SOUTHERN METHODIST UNIVERSITY
DALLAS
TEXAS
75222

1

DR. A. PETRASOVITS
ROOM 207B, FCCD AND DRUG ELDG.
TUNNEY'S PASTURE
OTTAWA, ONTARIO K1A-0L2,
CANADA

1

PROF. S. L. PFCENIX
SIBLEY SCHOOL OF MECHANICAL AND
AEROSPACE ENGINEERING
CORNELL UNIVERSITY
ITHACA
NY

1

14850

DR. A. L. POWELL
DIRECTOR
OFFICE OF NAVAL RESEARCH BRANCH OFF
495 SUMNER STREET
BOSTON
MA

1

02210

MR. F. R. PRIORI
CCCE 224 OPERATIONS TEST AND ONRS
EVALUATION FORCE (OPTVFOR)
NORFOLK
VIRGINIA
20360

1

PROF. M. L. PURI
DEPT. OF MATHEMATICS
P.O. BOX F
INDIANA UNIVERSITY FOUNDATION
BLOOMINGTON
IN

1

47401

PROF. H. ROBBINS
DEPARTMENT OF MATHEMATICS
COLUMBIA UNIVERSITY
NEW YORK,
NEW YORK. 10027

1

NO. OF COPIES

PROF. M ROSENBLATT
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF CALIFORNIA SAN DIEGO
LA JOLLA
CALIFORNIA

1

92093

PROF. S. M. ROSS
COLLEGE OF ENGINEERING
UNIVERSITY OF CALIFORNIA
BERKELEY
CA

1

94720

PROF. I RUBIN
SCHOOL OF ENGINEERING AND APPLIED
SCIENCE
UNIVERSITY OF CALIFORNIA
LOS ANGELES
CALIFORNIA 90024

1

PROF. J. R. SAVAGE
DEPARTMENT OF STATISTICS
YALE UNIVERSITY
NEW HAVEN,
CONNECTICUT
06520

1

PROF. L. L. JR. SCHARF
DEPARTMENT OF ELECTRICAL ENGINEERING
COLORADO STATE UNIVERSITY
FT. COLLINS,
COLORADO
80521

1

PROF. R. SERFLING
DEPARTMENT OF STATISTICS
FLORIDA STATE UNIVERSITY

1

TALLAHASSEE FLORIDA 32306

PROF. W. R. SCHUCANY
DEPARTMENT OF STATISTICS
SOUTHERN METHODIST UNIVERSITY
DALLAS,
TEXAS
75222

1

PROF. D. O. SIEGMUND
DEPT. OF STATISTICS
STANFORD UNIVERSITY
STANFORD
CA

1

94305

PROF. M. L. SPOOMAN
DEPT. OF ELECTRICAL ENGINEERING
POLYTECHNIC INSTITUTE OF NEW YORK
BROOKLYN,
NEW YORK
11201

1

PROF. N. SINGPURWALLA
DEPT. OF OPERATIONS RESEARCH
THE GEORGE WASHINGTON UNIVERSITY
707 22ND ST. N. W.
WASHINGTON, D. C.

1

20052

NO. OF COPIES

DR. A. I. SLAFKOSKY
SCIENTIFIC ADVISOR
COMMANCANT OF THE MARINE CORPS
WASHINGTON ,
C. C.
20380

1

MR. CHARLES S. SMITH
OASD (I&L),
PENTAGON
WASHINGTON
DC

1

20301

DR. D. E. SMITH
DESMATICS INC.
P.O. BOX 618
STATE COLLEGE
PENNSYLVANIA
16801

1

PROF. W. L. SMITH
DEPARTMENT OF STATISTICS
UNIVERSITY OF NORTH CAROLINA
CHAPEL HILL
NORTH CAROLINA 27514

1

PROF. H. SOLOMON
DEPARTMENT OF STATISTICS
STANFORD UNIVERSITY
STANFORD ,
CALIFORNIA
94305

1

MR. GLENN F. STAHLY
NATIONAL SECURITY AGENCY
FORT MEADE ,
MARYLAND 20755

1

MR. DAVID A. SWICK
ADVANCED PROJECTS GROUP
CODE 8103
NAVAL RESEARCH LAB.
WASHINGTON
DC

1

20375

MR. WENDELL G. SYKES
ARTHUR D. LITTLE, INC.
ACORN PARK
CAMBRIDGE
MA

1

02140

PROF. J. F. THOMPSON
DEPARTMENT OF MATHEMATICAL SCIENCE
RICE UNIVERSITY
HOUSTON,
TEXAS
77001

1

NO. OF COPIES

PROF. W. A. THOMPSON
DEPARTMENT OF STATISTICS
UNIVERSITY OF MISSOURI
COLUMBIA,
MISSOURI
65201

1

PROF. F. A. TILLMAN
DEPT. OF INDUSTRIAL ENGINEERING
KANSAS STATE UNIVERSITY
MANHATTAN
KS

1

66506

PROF. J. W. TUKEY
DEPARTMENT OF STATISTICS
PRINCETON UNIVERSITY
PRINCETON, N. J. 08540

1

PROF. A. F. VEINOTT
DEPARTMENT OF OPERATIONS RESEARCH
STANFORD UNIVERSITY
STANFORD
CALIFORNIA
94305

1

DANIEL H. WAGNER
STATION SQUARE ONE
FAULI, PENNSYLVANIA
19301

1

PROF. GRACE WAHBA
DEPT. OF STATISTICS
UNIVERSITY OF WISCONSIN
MADISON
WI

1

53706

PROF. K. T. WALLENIUS
DEPARTMENT OF MATHEMATICAL SCIENCES
CLEMSON UNIVERSITY
CLEMSON,
SOUTH CAROLINA 29631

1

PROF. G. S. WATSON
DEPARTMENT OF STATISTICS
PRINCETON, N. J. 08540

1

PROF. BERNARD WICROW
STANFORD ELECTRONICS LAB
STANFORD UNIVERSITY
STANFORD
CA

1

94305

PROF. G. E. WHITEHOUSE
DEPT. OF INDUSTRIAL ENGINEERING
LEHIGH UNIVERSITY
BETHLEHEM
PA

1

18015

NO. OF COPIES

PROF. S. ZACKS
DEPT. OF MATHEMATICS AND STATISTICS
CASE WESTERN RESERVE UNIVERSITY
CLEVELAND.,
OHIO
44106

1

PROF. N. ZIA-PASSAN
DEPARTMENT OF INDUSTRIAL AND
SYSTEMS ENGINEERING
ILLINOIS INSTITUTE OF TECHNOLOGY
CHICAGO
IL 60616

1

DR. DAVID BRILLINGER
STATISTICS DEPT.
UNIVERSITY OF CALIFORNIA
BERKELEY,
CALIFORNIA
94720

1

DR. CHIN CHEN
COMMONWEALTH ASSOCIATES, INC.
205 E. WASHINGTON
JACKSON, MI
49201

1

PROFESSOR PETER CHEN
ALFRED P. SLICAN SCHOOL OF MGMT
MASS INSTITUTE OF TECHNOLOGY
50 MEMORIAL DRIVE
CAMBRIDGE,
MASSACHUSETTS 02139

1

DR. E. CYNAR
O.R. DEPT.
NORTHWESTERN UNIVERSITY
EVANSTON,
ILLINOIS
60201

1

DR. GUY C. CORYNEN
L. 156 UNIV OF CALIFORNIA
LAWRENCE LIVERMORE LABORATORY
P. O. BOX 808
LIVERMORE, CALIFORNIA
94550

1

DR. MICHAEL A. CRANE
CONTROL ANALYSIS CORPORATION
800 WELCH ROAD
PALO ALTO, CALIFORNIA 94304

1

DR. D. F. CALEY
STATISTICS DEPT. (I A S)
AUSTRALIAN NATIONAL UNIVERSITY
CANBERRA, A. C. T. 2606
AUSTRALIA

1

NO. OF COPIES

HSU DER-ANN
SCH. OF ENGR./APPLIED SCI.
DEPT. OF CIVIL & GEOLOGICAL ENGR.
THE ENGINEERING QUADRANGLE
PRINCETON
NEW JERSEY 08540

1

ASSISTANT DEAN CAV ELDREDGE
SCHOOL OF BUSINESS ADMINISTRATION
UNIVERSITY OF EVANSVILLE
EVANSVILLE
INDIANA
47702

1

DR. G. W. FISHMAN
CURRICULUM IN O. R.
UNIVERSITY OF NORTH CAROLINA
PHILLIPS ANNEX, CHAPPEL HILL
NORTH CAROLINA
27514

1

DR. A. V. FOLLEN
DEPARTMENT OF PHYSIOLOGY
THE UNIVERSITY
LEEDS LS2 9JT
ENGLAND

1

DR. EUGENE C. HOMER
MANAGEMENT SCI. & ENGINEERING DEPT.
C.W. POST COLLEGE
OF LONG ISLAND UNIVERSITY
GREENVALE, NEW YORK
11548

1

DR. D. VERE JONES
DEPT. OF MATHEMATICS,
VICTORIA UNIVERSITY OF WELLINGTON,
P. O. BOX 196
WELLINGTON, NEW ZEALAND

1

J. M. LIITTSCHLAGER
SYSTEMS DIVISION
COLLEGE OF ENGINEERING
THE UNIVERSITY OF IOWA
IOWA CITY, IOWA
52242

1

Y. W. DE KWAADSTENIET
DEPARTMENT OF BIOLOGY
FREE UNIVERSITY
DE BOELELAAN 1087
AMSTERDAM-BUITENVELDER
THE NETHERLANDS

1

JACK P. LANCOLT
DEFENCE & CIVIL INST. OF ENVIR. MED
1133 SHEPARD AVE. WEST
PO BOX 2000 DOWNSVIEW
ONTARIO,
CANADA

1

NO. OF COPIES

MR. FRANK MCNOLTY
1142 PCME AVE.
SUNNYVALE, CA
94087

1

DR. P. A. F. MCGRAN
STATISTICS DEPT. (I A S)
AUSTRALIAN NATIONAL UNIVERSITY
CANBERRA, A. C. T 2606
AUSTRALIA

1

G. W. MCRRISON
OAK RIDGE NATIONAL LABORATORY
OAK RIDGE, TENNESSE
37830

1

MR. BRADLEY NICKY
USCA - FOREST SERVICE
P. O. BOX 5007
FIVERSICE
CALIFORNIA
92507

1

DR. DAVID CATES
DEPT. OF STATISTICS
HARVARD UNIVERSITY
SCIENCE CENTER, 1 OXFORD ST.
CAMBRIDGE, MA.
02132

1

Professor Emanuel Parzen, Director
Institute of Statistics.
Texas A&M University
College Station, TX 77843

1

MR ART PETERSON
DEPT. OF BIOSTATISTICS SC-32
UNIVERSITY OF WASHINGTON
SEATTLE, WASHINGTON
98195

1

DAVID W. ROBINSON
CONTROL ANALYSIS CORPORATION
800 WELCH ROAD
PALO ALTO
CALIFORNIA
94304

1

C. H. SAUER (21-129)
THOMAS J. WATSON RESEARCH CENTER
P.O. BOX 218
YORKTOWN HEIGHTS, NEW YORK
10598

1

NO. OF COPIES

MR. C. A. STEELE JR
P.O. BOX 45
MAGNOLIA, MA
01930

1

FRCE P. SULLO
SCHOOL OF MANAGEMENT
RENSSELAER POLYTECHNIC
INSTITUTE
TROY, N.Y.
12181

1

DR. ROBERT TRAPPL
INSTITUT FUR ALLGEMEINE UND
VERGLEICHENDE PHYSIOLOGIE DER
UNIVERSITAT WIEN
SCHWARZSPANIERSTRASSE 17
A-1090 WIEN OSTERREICH (AUSTRIA)

1

PALL A. WILLIS
POLY TECHNICA
2824 W. GEORGE MASON
FALLS CHURCH, VA. 22042

1

TUNCEL M. YEGULALP
KRLMB SCHOOL OF MINES
918 S.W. MIDD
COLUMBIA UNIVERSITY
520 W. 12TH ST.
NEW YORK, N.Y. 10027

1

SHIEGKI YOKOYAMA
RESEARCH CENTER OF APPLIED
INFORMATION SCIENCES
TOHOKU UNIVERSITY
KATAHIRA-CHO 2-1,
SENDAI 980 JAPAN

1

Dean of Research
Code 012
Naval Postgraduate School
Monterey, Ca. 93940

1

Library (Code 55)
Naval Postgraduate School
Monterey, CA 93940

1

NO. OF COPIES

Professor Peter Lewis
Code 55Lw
Naval Postgraduate School
Monterey, Ca. 93949

50

Mr. G. Shedler
K53-282
IBM Research Lab
Cottle Road
San Jose, Ca. 95193

50

R. J. Stampfel
Code 55
Naval Postgraduate School
Monterey, Ca. 93940

1